

Landsteini Van der Scheer

J - Exp. Med 1936

63, p. 325

(rejection crises)

TABLE XXXIV

MOLECULAR COMPOSITION OF SPECIFIC PRECIPITATES

Antigen	Antibody	Empirical composition of precipitate at				Composition of soluble compound in zone of partial inhibition
		Extreme ^a antibody excess	Antibody excess end of equivalence zone	Antigen excess end of equivalence zone	Zone of partial inhibition	
Ovalbumin	Rabbit	A ₅ G	A ₃ G	A ₅ G ₂	A ₂ G	(AG)
Dye-ovalbumin	Rabbit	(A ₅ G)	(A ₃ G)	A ₅ G ₂	A ₃ G ₄	(AG ₂)
Serum albumin	Rabbit	A ₆ G	A ₄ G	A ₃ G	A ₂ G	(AG)
Thyroglobulin	Rabbit	A ₄₀ G	A ₁₄ G	A ₁₀ G	A ₂ G	(AG)
Viviparus hemocyanin	Rabbit	—	A ₁₂₀ G	A ₈₃ G	A ₃₅ G	—
Tobacco mosaic virus	Rabbit	A ₉₀₀ G	A ₄₅₀ G	—	—	—
Diphtheria toxin	Horse	A ₈ G	A ₄ G	A ₂ G ₂	AG	(AG ₂)
Ovalbumin	Horse	(A ₄ G)	A ₂ G	—	AG	(AG ₂)

A = antibody, G = antigen.

Formulas in parentheses are somewhat uncertain.

^a For meaning of this and similar terms, see Figures 31a and 31b.