

## Invasive Bladder Testing Before Incontinence Surgery May Be Unnecessary

May 02, 2012 |

**N** *IH-Funded Study Finds Office Visit Alone Just as Effective*

Invasive and costly tests commonly performed on women before surgery for stress urinary incontinence (SUI) may not be necessary, according to researchers at the University of California San Diego, School of Medicine and the Urinary Incontinence Treatment Network. The study, supported by the National Institutes of Health (NIH), will be released online May 2 by the *New England Journal of Medicine (NEJM)*.



Nager discusses study findings.

The study, which compared results after a combination of a preoperative check-up and bladder function tests to an office check-up alone, found that women who only had the office examination experienced equally successful outcomes after surgery.

“We wanted to know if invasive tests are really needed in women who have SUI, or if observation alone could achieve the same outcomes,” said study lead -author Charles Nager, MD, director of Urogynecology and Reconstructive Pelvic Surgery at UC San Diego Health System. “The findings of our study argue that the tests provide no added benefit for surgical treatment success to patients.”

SUI affects up to 30 million American women and causes leakage of urine when coughing, laughing, sneezing, running or lifting heavy objects. Bladder tests that use technology such as

imaging, pressure monitors and muscle and nerve sensors, are commonly used on these women before surgery to characterize the degree of incontinence and to guide decisions about treatment options. However, the tests are uncomfortable, costly and increase the risk of urinary tract infections.

In the study, half of a group of 630 women with uncomplicated SUI underwent a preoperative office evaluation with bladder function testing. The other half had an office evaluation only. The proportion of women who achieved treatment success was similar, 76.9 percent versus 77.2 percent, respectively, with no significant differences between groups in quality of life, patient satisfaction or voiding dysfunction – an inability to empty the bladder completely.

UC San Diego School of Medicine researchers who participated in this study included Michael Albo, MD, and Emily Lukacz, MD.

The study was funded by the National Institute of Diabetes and Digestive and Kidney Diseases and the Eunice Kennedy Shriver National Institute of Child Health and Human Development.

The Women's Pelvic Medicine Center at UC San Diego Health System is the region's first center designed exclusively for the diagnosis and treatment of pelvic floor disorders. Since the program's inception in 1998, women with complex conditions, such as incontinence and pelvic organ prolapse, have been treated through advanced medical therapies and minimally invasive surgical techniques.

# # #

Media Contact: Michelle Brubaker, 619-543-6163, [mubruber@ucsd.edu](mailto:mubruber@ucsd.edu)

---

## Related Specialties

---

[Women's Pelvic Medicine](#)

## Share This Article

---



## Related News

---

[Young, Healthy Woman Suffers  
Brain Inflammation after Mild  
COVID-19 Infection](#)  
8/9/2021

Organ Transplant Recipients  
Significantly Protected by COVID-19  
Vaccination  
8/5/2021

Rethinking Remdesivir  
8/2/2021

Obesity and Cardiovascular Factors  
Combine to Cause Cognitive  
Decline in Latinos  
7/28/2021

[View All News >](#)

## Follow Us

---

 Follow @ucsdhealth

