

Learning Impacts How the Brain Processes What We See

July 13, 2015 | Yadira Galindo

From the smell of flowers to the taste of wine, our perception is strongly influenced by prior knowledge and expectations, a cognitive process known as top-down control.

In a University of California, San Diego School of Medicine study published July 13 in the online journal *Nature Neuroscience*, a research team led by Takaki Komiyama, PhD, assistant professor of neurosciences and neurobiology, reports that in mouse models, the brain significantly changed its visual cortex operation modes by implementing top-down processes during learning.

“We found that when the mouse assigns a new meaning to a previously neutral visual stimulus, top-down control becomes much more influential in activating the visual cortex,” said first author Hiroshi Makino, PhD, postdoctoral researcher in Komiyama’s lab. “Top-down inputs interact with specific neuron types in the visual cortex to modulate its operation modes.”

This cognitive process uses our thoughts and influences our senses. For example, when we see a word with missing letters, our brain is able to fill in the blank based on past experiences.

Researchers looked at activity in excitatory neurons and somatostatin-expressing inhibitory neurons in the visual cortex and top-down inputs from the retrosplenial cortex (RSC) during associative learning to see how these affected the top-down and bottom-up processing—when perception begins with the senses.

The findings indicate that intricate interactions of various circuit components effectively change the balance of top-down and bottom-up processing, with learning enhancing the contribution of top-down control. These results support the long-standing theory that the brain does not faithfully represent the environment but rather attempts to predict it based upon prior information.

“In addition to revealing circuit mechanisms underlying these learning-related changes, our findings may have implications in understanding the pathophysiology of psychiatric diseases, such as schizophrenia, that generate abnormal perception,” said Makino.

Funding for this research came, in part, from the National Institutes of Health (1R01NS091010-01, 1R01DC014690-01), Japan Science and Technology Agency (PRESTO), Pew Charitable Trusts, Alfred P. Sloan Foundation, David & Lucile Packard Foundation, Human Frontier Science Program, McKnight Foundation and New York Stem Cell Foundation.

Care at UC San Diego Health

Psychiatry and Behavioral Health

Media Contact

Yadira Galindo
858-249-0456
ygalindo@health.ucsd.edu

Share This Article



Related News

[Researchers Identify Specific Genetic Vulnerabilities to PTSD among U.S. Veterans](#)
7/29/2019

[Study to Explore Whether Cannabis Compound Eases Severe Symptoms of Autism](#)
4/26/2018

[Negative Fateful Life Events and the Brains of Middle-Aged Men](#)
4/5/2018

[Genetic Assessment Developed to Determine Risk for Age-Associated Alzheimer's Disease](#)
3/21/2017

[View All News >](#)

Follow Us

 Follow [@ucsdhealth](#)

