


Autism Genes Activate During Fetal Brain Development

Mutations that cause autism in children are connected to a pathway that regulates brain development.

February 18, 2015 | Scott LaFee

Scientists at the University of California, San Diego School of Medicine have found that mutations that cause autism in children are connected to a pathway that regulates brain development. The research, led by Lilia Iakoucheva, PhD, assistant professor in the Department of Psychiatry, is published in the February 18 issue of *Neuron*.

The researchers studied a set of well-known autism mutations called copy number variants or CNVs. They investigated when and where the genes were expressed during brain development. "One surprising thing that we immediately observed was that different CNVs seemed to be turned on in different developmental periods," said Iakoucheva.



Autism mutations may influence brain size through RhoA pathway during fetal brain development.

Specifically, the scientists noted that one CNV located in a region of the genome known as 16p11.2, contained genes active during the late mid-fetal period. Ultimately, they identified a network of genes that showed a similar pattern of activation including KCTD13 within 16p11.2 and CUL3, a gene from a different chromosome that is also mutated in children with autism.

"The most exciting moment for us was when we realized that the proteins encoded by these genes form a complex that regulates the levels of a third protein, RhoA," said Iakoucheva. Rho proteins play critical roles in neuronal migration and brain morphogenesis at early stages of brain development. "Suddenly, everything came together and made sense."

Further experiments confirmed that CUL3 mutations disrupt interaction with KCTD13, suggesting that 16p11.2 CNV and CUL3 may act via the same RhoA pathway. RhoA levels influence head and body size in zebrafish, a model organism used by geneticists to investigate gene functions.

Children with 16p11.2 CNV also have enlarged or decreased head sizes and suffer from obesity or are underweight. "Our model fits perfectly with what we observe in the patients," said Guan Ning Lin, PhD, a fellow in Iakoucheva's laboratory and co-first author with Roser Corominas, PhD.

Interestingly, the RhoA pathway has recently been implicated in a rare form of autism called Timothy syndrome, which is caused by the mutation in a completely different gene. "The fact that three different types of mutations may act via the same pathway is remarkable," said Iakoucheva. "My hope is that we would be able to target it therapeutically."

Iakoucheva and colleagues are planning to test RhoA pathway inhibitors using a stem cell model of autism. "If we can discover the precise mechanism and develop targeted treatments for a handful of children, or even for a single child with autism, I would be happy," she said.

Co-authors include Xiping Yang, David E. Hill and Marc Vidal, Dana-Farber Cancer Institute; Irma Lemmens and Jan Tavernier, Ghent University, Belgium; and Jonathan Sebat, Beyster Center for Genomics of Psychiatric Diseases and UCSD.

This research was funded, in part, by National Institutes of Health (grants R01MH091350, R01HD065288, R21MH104766 and R01MH105524).

Care at UC San Diego Health System

[Autism Center of Excellence](#)

[Psychiatry and Behavioral Health](#)

Media Contact

Scott LaFee
858-249-0456
slafee@health.ucsd.edu

Share This Article



Related News

Predicting Autism Risk May Begin

With a Drop of Blood

2/13/2020

Study to Explore Whether Cannabis Compound Eases Severe Symptoms of Autism

4/26/2018

Inflamed Support Cells Appear to Contribute to Some Kinds of Autism

10/18/2017

Mini-brain Model of Idiopathic Autism Reveals Underlying Pathology of Neuronal Overgrowth

7/13/2016

[View All News >](#)

Follow Us

 Follow @ucsdhealth

