CHEMOSTAT


AIR
CAPILLARY LEAK
INOCULATING PORT
AIR ESCAPE
PRESSURE REGULATOR
AIR
LEVELING SIPHON
COLLECTING VESSEL
GROWTH TUBE
SINTERED DISK
RESERVOIR

Caffeine	19
Theophylline	11
Paraxanthine	8.4
Theobromine	7.5
Tetramethyluric acid	7.0
8-Methoxycaffeine	5.2
8-Chlorocaffeine	4.2
Azaguanine	3.4
Benzimidazole	3.0
Adenine	2.8


2000
1500
1000
500
T5 MUTANTS PER CC.
50
100
150
200
250
300
350
400
450
GENERATIONS ($\frac{t}{\tau}$)


2.0
1.5
1.0
0.5
OPTICAL DENSITY (280mμ)
4
8
12
τ (HOURS)


1.0
0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
OPTICAL DENSITY
SUPERNATANT OBSERVED AT 280 mμ
SUPERNATANT CORRECTED
BACTERIA AT 350 mμ
0
0.5
1.0
1.5
2.0
2.5
TIME (HOURS)

B/r/lt

	Aerobic		Anaerobic	
	T5	T6	T5	T6
Spontaneous	4.0	1.0	0.6	(0.4)
Adenosine	1.2	0.4	0.8	0.6
Theophylline	22	2.1	1.1	1.3
Theophylline + Adenosine	1.4	0.4	--	--
U - V	15	5.6	13.6	7.7
X-Ray	16.4	11.5	7.0	5.4
Spontaneous High Growth Rate (B/lt)	3.8	3.0	3.8	3.0



3.0-164

ADENINE



GUANINE



2.0-500

HYPOXANTHINE



XANTHINE



17-150

CAFFEINE



10.8-150

THEOPHYLLINE



7.5-150

THEOBROMINE



8.4-150

PARAXANTHINE



7.6-150

TETRAMETHYL URIC ACID



4.3-150

8-CHLOROCAFFEINE



5.2-150

8-METHOXYCAFFEINE



3.4-150

8-AZAGUANINE

	T 5 Resistance	T 6 Resistance
Spontaneous	4.0	1.0
150 mg/l Adenosine	1.2	0.4


MUTATION RATE (X 10^8) PER HOUR
20
10
200
400
600
800
THEOPHYLLINE CONCENTRATION (mg/l)


B/r/I,t
T5 RESISTANCE
T6 RESISTANCE
MUTATION RATE (PER 10^8 BACTERIA/hr.)
30
20
10
5
10
15
20
U-V INTENSITY (ergs/mm^2/hour)


FRACTION OF ARGININE CONTENT TAKEN FROM OUTSIDE
1.0
0.8
0.6
0.4
0.2
0
0.2
0.4
0.6
ARGININE CONCENTRATION (γ/ℓ)


τ = 8 HRS.
1150 γ/ℓ ARGININE (0.5)
690 γ/ℓ ARGININE (0.3)
230 γ/ℓ ARGININE (0.1)
0 γ/ℓ ARGININE (0)
ARGININELESS BACTERIA PER CC.
10^7
10^6
10^5
10^4
20
40
60
HOURS


B/l, t
τ = 5.5 hr
T5 RESISTANT MUTANTS PER 10^8 BACTERIA
600
400
200
8 1/2 hr
150 mg/ℓ CAFFEINE
ADDED AT 68 hr
25
50
75
100
120
TIME (HOURS)


10^8
SLOW
TOTAL
BACTERIA
BACTERIA / mℓ
FAST
0
25
50
75
150
T 5
RISISTANT
MUTANTS
MUTANTS / mℓ
100
50
0
25
50
75
1.5
TRYPTOPHAN
CONCENTRATION
1
0.5
μg/ℓ TRYPTOPHAN
0
25
50
75
GENERATIONS


T5 RESISTANTS PER 10^8 BACTERIA
300
200
100
25
50
75
100
GENERATIONS


TRYPTOPHANLESS — HISTIDINELESS
τ = 6.4 hr.
AT 280 mμ
AT 250 mμ
COMPUTED COMPOUND AT 280 mμ
$e^{-\frac{t}{7\,hr.}}$
OPTICAL DENSITY
0.6
0.4
0.2
5
10
15
20
25
30
HOURS


0.6
0.4
0.2
GROWTH RATE (hr⁻¹)
ARGININELESS B (D84/6)
1
2
3
4
5
ARGININE CONCENTRATION (g/l)


1.25
1.00
0.75
0.50
0.25
0.0
OPTICAL DENSITY
240
260
280
300
320
WAVELENGTH (mμ)


MUTANTS PER 10^8 BACTERIA
1500
1000
500
CAFFEINE ~ 19 X 10^{-8}/HR
THEOPHYLLINE ~ 10.8 X 10^{-8}/HR
PARAXANTHINE ~ 8.4 X 10^{-8}/HR
THEOBROMINE ~ 7.5 X 10^{-8}/HR
8-AZAGUANINE ~ 3.4 X 10^{-8}/HR
50
100
150
200
HOURS


MUTANTS PER 10^8 BACTERIA
150
100
50
TRYPTOPHANE 1.55 X 10^{-8}/HR
NITROGEN 0.82 X 10^{-8}/HR
PHOSPHORUS 0.67 X 10^{-8}/HR
LACTATE 0.53 X 10^{-8}/HR
SPONTANEOUS MUTATIONS
50
100
150
200
HOURS


T5 RESISTANT MUTANTS PER CC
2000
1500
1000
500
600 mg/l
150 mg/l
50 mg/l
15 mg/l
NO THEOPHYLLINE
20
40
60
80
100
120
HOURS


T5 RESISTANT MUTANTS PER CC
1500
1000
500
τ = 12 hr
τ = 6 hr
τ = 2 hr
10
20
30
GENERATIONS (t/τ)

Hypoxanthine
Xanthine
2,6 Diaminopurine
8 Methylxanthine
Imidazole

Uracil
Thymine
6-Methyluracil
5-Bromouracil
5-Aminouracil
3-Methyluracil
Barbituric Acid

Colchicine
Urethane
Paraldehyde
Allantoin
Acetysalicylic acid
Procaine
Quinine sulfate
Aminopyrine
Atropine


α_f
0.8
FAST
0.6
α_o
GROWTH RATE α (hr.$^{-1}$)
0.4
SLOW
0.2
α_s
C_2
C_1
1
2
3
4
TRYPTOPHAN CONCENTRATION (μg/ℓ)


0.8
0.6
0.4
0.2
GROWTH RATE α (hr⁻¹)
FAST
SLOW
1
2
3
4
5
TRYPTOPHANE CONCENTRATION (γ/l)


OPTICAL DENSITY
1.6
1.2
0.8
0.4
280 mμ
250 mμ
0
2
4
6
8
HOURS


T5 RESISTANT MUTANTS PER 10^8 BACTERIA
150
100
50
0
A
B
0
30
60
90
120
HOURS


Mutants per 10^8 Bacteria
80
60
40
20
0
40
80
120
160
200
240
HOURS
$\tau = 2hr$
$\frac{\lambda}{\tau} = 0.30 \times 10^{-8}/hr$
$\tau = 5hr$
$\frac{\lambda}{\tau} = 0.30 \times 10^{-8}/hr$
$\tau = 10hr$
$\frac{\lambda}{\tau} = 0.29 \times 10^{-8}/hr$
T6 Resistance
Tryptophane Control



FIGURE 7. A projection of two successive residues of *one* chain of the structure. The direction of projection is parallel to the fibre axis. The figures show the height of each atom (in ångströms) above the level of the lower base.

TABLE 1. CO-ORDINATES FOR THE ATOMS OF THE BACKBONE, FOR A SINGLE RESIDUE

atom	ρ (Å)	ϕ	Z (Å)
P	10·0	0·0°	0·0
O_{I}	8·95	− 3·6°	+0·8
O_{II}	11·25	+ 0·7°	+0·8
O_{III}	9·65	+ 8·9°	−0·5
O_{IV}	10·35	− 5·3°	−1·3
$C_{5'}$	9·6	−22·2°	−2·8
$C_{4'}$	9·65	−13·2°	−3·2
$C_{3'}$	9·2	− 7·3°	−2·05
$C_{2'}$	8·65	+ 0·4°	−2·8
$C_{1'}$	8·2	− 3·5°	−4·15
$O_{1'}$	8·8	−11·8°	−4·35
N	6·7	− 4·2°	−4·15
diad	—	+39·0°	−4·15

$$R.CH(NH_2).COOH.$$

$$NH_2\overset{\overset{\displaystyle R}{|}}{C}H.CO\left[-NH.\overset{\overset{\displaystyle R}{|}}{C}H.CO-\right]_n NH.\overset{\overset{\displaystyle R}{|}}{C}H.COOH.$$



Structure of cattle insulin.

TABLE II

The amino acid sequence in positions A7–10 of insulins from various species

Cattle	..	..	$CySO_3H$.Ala.Ser.Val
Pig	..	..	$CySO_3H$.Thr.Ser.Ileu
Sheep	..	..	$CySO_3H$.Ala.Gly.Val
Horse	..	..	$CySO_3H$.Thr.Gly.Ileu
Whale	..	..	$CySO_3H$.Thr.Ser.Ileu

TABLE I

Common naturally occurring amino acids

Amino acid	*Structure of R group*	*Abbreviation used for residue —NH—CH(R)—CO—*	*Number of residues in insulin*
Aliphatic			
Glycine	—H	Gly	4
Alanine	$-CH_3$	Ala	3
Valine	$-CH(CH_3)_2$	Val	5
Leucine	$-CH_2-CH(CH_3)_2$	Leu	6
Isoleucine	$-CH(CH_3)-CH_2-CH_3$	Ileu	1
Serine	$-CH_2-OH$	Ser	3
Threonine	$-CH(OH)-CH_3$	Thr	1
Aromatic			
Phenylalanine	$-CH_2-C_6H_5$	Phe	3
Tyrosine	$-CH_2-C_6H_4-OH$	Tyr	4
Tryptophan	$-CH_2-C_8H_6N$	Try	0
Acidic			
Aspartic acid	$-CH_2-COOH$	Asp	0
Glutamic acid	$-CH_2-CH_2-COOH$	Glu	4
Acid amides			
Asparagine	$-CH_2-CONH_2$	$AspNH_2$	3
Glutamine	$-CH_2-CH_2-CONH_2$	$GluNH_2$	3
Basic			
Lysine	$-CH_2-CH_2-CH_2-CH_2-NH_2$	Lys	1
Arginine	$-CH_2-CH_2-CH_2-NH-C(=NH)-NH_2$	Arg	1
Histidine	$-CH_2-C_3H_3N_2$	His	2
S-containing			
Cysteine	$-CH_2-SH$	CySH	0
Cystine	$-CH_2-S-S-CH_2-$	CyS—CyS	3
Methionine	$-CH_2-CH_2-S-CH_3$	Met	0
Cyclized			
Proline	H_2C-CH_2*; H_2C, $CH-COOH$, N, H	Pro	1

* Complete structure of the amino acid.

SUPPRESSOR GENE

td MUTANT		su_2	su_3	su_{24}	su_6
Type I	td_1	−	−	−	−
Type II	td_2	+	−	−	+
Type III	td_3	−	+	+	−
	td_{24}	−	+	+	−
Type IV	td_6	−	−	−	+

+ = SUPPRESSION



FRAGMENT OF CHAIN of desoxyribonucleic acid shows the three basic units that make up the molecule. Repeated over and over in a long chain, they make it 1,000 times as long as it is thick. The backbone is made up of pentose sugar molecules (marked by the middle colored square), linked by phosphate groups (bottom square). The bases (top square), adenine, cytosine, guanine and thymine protrude off each sugar in irregular order.



ONE LINKAGE of base to base across the pair of DNA chains is between adenine and thymine. For the structure proposed, the link of a large base with a small one is required to fit chains together.



ANOTHER LINKAGE is comprised of guanine with cytosine. Assuming the existence of hydrogen bonds between the bases, these two pairings, and only these, will explain the actual configuration.





STRUCTURAL MODEL shows a pair of DNA chains wound as a helix about the fiber axis. The pentose sugars can be plainly seen. From every one on each chain protrudes a base, linked to an opposing one at the same level by a hydrogen bond. These base-to-base links act as horizontal supports, holding the chains together. Upper photograph is a top view.


ASPARAGINE GROWN
L-TRYPTOPHAN GROWN
OXYGEN UPTAKE MICROLITERS
300
200
100
0
L-TRYPTOPHAN
ENDOGENOUS
L-TRYPTOPHAN
ENDOGENOUS
0
80
160
0
80
160
MINUTES

$$\text{Gal}^* + \text{ATP} \xrightarrow{\text{Galactokinase}} \text{Gal}^*\text{-1-P} + \text{ADP} \qquad (1)$$

$$\text{Gal}^*\text{-1-P} + \text{UDPG} \underset{}{\overset{\text{Gal-1-P uridyl transferase}}{\rightleftharpoons}} \text{UDPGal}^* + \text{G-1-P} \qquad (2)$$

$$\text{UDPGal}^* \underset{\text{UDPGal-4-epimerase}}{\rightleftharpoons} \text{UDPG}^* \qquad (3)$$

The sum of reactions 1, 2, and 3 is

$$\text{Gal} + \text{ATP} \longrightarrow \text{G-1-P} + \text{ADP}$$



Fig. 2. The biochemical pathway for the oxidation of L-tryptophan by *Pseudomonas fluorescens*. The following inducible enzymes are involved: (a) tryptophan peroxidase + oxidase; (b) formylase; (c) kynureninase; (d) anthranilic acid oxidase (probably a multienzyme system); (e) pyrocatechase; (f) lactonizing enzyme; (g) delactonizing enzyme. The further breakdown of β-ketoadipic is catalyzed by an inducible enzyme, the decomposition of alanine is a constitutive reaction.



Fig. 53. Biosynthesis of the aromatic amino acids. Known positions of genetic interference in reaction series are indicated by the circled numbers on the cross bars. (See text for full explanation.)



Fig. 59. Biosynthesis of the purines of nucleic acids. Genetic blocks indicated by the cross bars and circled numbers are discussed in detail in the text.



Fig. 57. The biosynthesis of lysine in *Neurospora* and *E. coli*. Two distinct pathways for biosynthesis are indicated.



Fig. 54. Biosynthesis of the sulfur amino acids and threonine. The cross bars indicate positions of known genetic blocks.



Fig. 58. The biosynthesis of leucine, isoleucine, and valine. It has been shown that a single mutation can block similar reactions in two series as indicated by the cross bar just following the keto acids that correspond to isoleucine and valine.



Fig. 55. The biosynthesis and degradation of histidine. Circled numbers indicate different mutant types known in *Neurospora* or *E. coli*.



Fig. 60. Biosynthesis of some of the pyrimidines of nucleic acids. Studies of mutants have not established whether orotic acid and orotidine are true intermediates, but enzyme work has shown that orotic acid goes to orotidine phosphate and thence to the phosphates of uridine and cytidine instead of to the free nucleosides as indicated here.



Fig. 56. Arginine and proline biosynthesis. Cross bars indicate apparent positions of known genetic blocks.

RESPONSES TO FIRST, SECOND AND THIRD INJECTIONS OF I'BGG
RABBIT # 977



RESPONSES TO FIRST, SECOND AND THIRD INJECTIONS OF I'BSA
RABBIT # 984




Paragene
Ntd
E
M



Paragene


Add 2×10⁻³ TMG to tank
2x10⁻⁴ TMG
Strain M L3 Sept 57
Dried limited ≈ 2.0 mg/L NH4Cl
pH 6.65
gas phase: Air
τ = 2.95 hrs
S.A.
0.25
0.2
0.15
0.1
0.05
0
1
2
3
4
5
6
7
Generations


$\frac{N_d}{10^3} = 10$
$\bar{N}$
$\bar{N} = \frac{1 - e^{-N_d h}}{e^h - 1}$
$N_d = 10^4$
$\bar{n} = \frac{1}{h}$
0
1
$\ln \bar{n}$
10^1


N_d
$\bar{N} = \frac{1 - e^{-N_d h}}{e^h - 1}$
$N_d = 10^4$
$\bar{n} = \frac{1}{h}$
$\bar{N}$
0
10^3
$\ln \bar{n}$
10^4


mandelic acid
benzyl alcohol
CHO
COOH
$\frac{1}{2}$ O_2
OH
$\frac{1}{2}$ O_2
COOH
NH_2
tryptophane
anthranilic acid
$\frac{1}{2}$ O_2
$(-CO_2)$
$(-NH_3)$
O_2
$-CO_2$
OH
OH
COOH
P-OH-benzoic acid
OH
O_2 (?via quinone?)
$\frac{1}{2}$ O_2
COOH
COOH
CH
CH
COOH
CH
CH
cis-cis-muconic acid
protocatechuic acid
HO
OH
O_2
$(-CO_2)$
β-ketoadipic acid
COOH
CH_2
COOH
CO
CH_2
CH_2
COOH
α-ketoglutaric acid
CO
CH_2—CH_2COOH


Precursor in Neurospora
(1)
NH2(CH2)3CHCOOH
NH2
ornithine
+ CO2 + NH3
(2)
O=C—NH(CH2)3CHCOOH
NH2
NH2
citrulline
+ NH3
(3)
NH=C—NH(CH2)3CHCOOH
NH2
NH2
arginine
+ H2O


1.0
0.8
0.6
0.4
0.2
0
Fraction of Maximum Activity
0
0.5
1.0
1.5
2.0
Generations


TMG = 10^-4 M τ = 3.12 hrs
τ = 9.1 hrs
τ = 3 hrs
τ = 3.1 hrs 4×10^-6 TMG
Strain M. in Dance med
20 mg/L NH4Cl pH 6.7
after 642-644
S.A.
2
1.5
1.0
0.8
0.6
0.4
0.2
0
1
2
3
4
5
6
Generations


.06
.04
.02
0
Fraction of Maximum Activity
Saturation
S
0
2
4
6
8
10
12
Generations


VALVE
CLOCK
LEAK
PORT
LEAK
AIR
STORAGE TANK
PORT
AIR ESCAPE
AIR
h_1
h_2
h_3
h_4
GROWTH TUBE
COLLECTING VESSEL
PRESSURE REGULATORS

$$\text{Acetyl-ornithine} \xrightarrow[G(-2)]{E(-2)} \text{Ornithine} \xrightarrow[G(-1)]{E(-1)} \text{Citrulline} \xrightarrow[G(0)]{E(0)} \text{Arginine}$$


Paragene
Head
"Enzyme"
Ntd
M


T5 RESISTANT MUTANTS PER 10^8 BACTERIA
900
600
300
0
C
D
GUANOSINE ADDED
0
50
100
150
200
HOURS